



TITLE:

Single-molecule observations of RNA-RNA kissing interactions in a DNA nanostructure

AUTHOR(S):

Takeuchi, Yosuke; Endo, Masayuki; Suzuki, Yuki; Hidaka, Kumi; Durand, Guillaume; Dausse, Eric; Toulmé, Jean Jacques; Sugiyama, Hiroshi

CITATION:

Takeuchi, Yosuke ...[et al]. Single-molecule observations of RNA-RNA kissing interactions in a DNA nanostructure. Biomaterials Science 2016, 4: 130-135

ISSUE DATE:

2016-01-01

URL:

<http://hdl.handle.net/2433/218508>

RIGHT:

© 2015 The Royal Society of Chemistry.



Cite this: *Biomater. Sci.*, 2016, **4**, 130

Single-molecule observations of RNA–RNA kissing interactions in a DNA nanostructure†

Yosuke Takeuchi,^a Masayuki Endo,^{*b} Yuki Suzuki,^a Kumi Hidaka,^a Guillaume Durand,^{c,d} Eric Dausse,^{c,d} Jean-Jacques Toulmé^{*c,d} and Hiroshi Sugiyama^{*a,b}

RNA molecules uniquely form a complex through specific hairpin loops, called a kissing complex. The kissing complex is widely investigated and used for the construction of RNA nanostructures. Molecular switches have also been created by combining a kissing loop and a ligand-binding aptamer to control the interactions of RNA molecules. In this study, we incorporated two kinds of RNA molecules into a DNA origami structure and used atomic force microscopy to observe their ligand-responsive interactions at the single-molecule level. We used a designed RNA aptamer called GTPswitch, which has a guanosine triphosphate (GTP) responsive domain and can bind to the target RNA hairpin named Aptakiss in the presence of GTP. We observed shape changes of the DNA/RNA strands in the DNA origami, which are induced by the GTPswitch, into two different shapes in the absence and presence of GTP, respectively. We also found that the switching function in the nanospace could be improved by using a cover strand over the kissing loop of the GTPswitch or by deleting one base from this kissing loop. These newly designed ligand-responsive aptamers can be used for the controlled assembly of the various DNA and RNA nanostructures.

Received 25th July 2015,
Accepted 25th September 2015

DOI: 10.1039/c5bm00274e

www.rsc.org/biomaterialsscience

Introduction

Structural diversity of RNA is one of the important properties of RNA molecules, which exhibit unique functions such as specific complex formation and catalysis. One of the variations of the complex formation between RNA molecules includes a “kissing complex,” which enables assembly of complementary RNA loops *via* Watson–Crick base pairing.^{1–8} This specific kissing complex formation with native RNA motifs has been used to create various RNA architectures such as polygonal structures and three-dimensional assemblies.^{9–11} Aptamers with a kissing RNA loop have been artificially developed by *in vitro* selection.^{12–15} A ligand-responsive kissing aptamer, called a guanosine triphosphate switch (GTPswitch), has been recently reported.¹⁶ This GTPswitch has a GTP-binding domain and a kissing domain that binds to a target RNA loop (Aptakiss) in the presence of GTP.¹⁶ Herein, we tried to visualize this unique

ligand-responsive switching interaction between the GTPswitch and its counterpart Aptakiss at single-molecule resolution.

Direct observation of interactions between biomolecules by using atomic force microscopy (AFM) is one of the practical methods for characterizing the properties of complex formation.¹⁷ Especially using a high-speed AFM and nano-sized DNA origami scaffold called DNA frame,^{17,18} dynamic formation of G-quadruplex structure and double-stranded DNA has been visualized.^{19–22} This single-molecule observation system should be used to observe the interactions of kissing aptamers for characterizing their properties.

In this study, we placed the GTPswitch and the target kissing loop Aptakiss into the cavity of a DNA frame and directly observed their interaction (Fig. 1a).^{19–22} The GTPswitch was generated based on a KG51 RNA kissing hairpin, which can bind to the Aptakiss only in the presence of GTP.¹⁶ We elongated the 5′ end of each RNA molecule and hybridized these molecules to the supporting DNA strands. These constructs were incorporated into the DNA frame through four ssDNA linkers (Fig. 1b). When the GTPswitch binds to the Aptakiss upon the addition of GTP, the configuration change of the supporting DNA strands from the unbound “double-loop” to the “X-shape” should be observed in the DNA frame (Fig. 1c). We investigated the ligand-responsive binding of the GTPswitch to the Aptakiss by observing the structural changes in the supporting DNA strands in the DNA frame, and examined the experimental conditions to improve the switching function.

^aDepartment of Chemistry, Graduate School of Science, Kyoto University, Kitashirakawa-oiwakecho, Sakyo-ku, Kyoto 606-8502, Japan

^bInstitute for Integrated Cell-Material Sciences (WPI-iCeMS), Kyoto University, Yoshida-ushinomiyacho, Sakyo-ku, Kyoto, 606-8501, Japan.

E-mail: endo@kuchem.kyoto-u.ac.jp, hs@kuchem.kyoto-u.ac.jp

^cARNA laboratory, University of Bordeaux, 146 rue Léo Saignat, 33076 Bordeaux, France. E-mail: jean-jacques.toulme@inserm.fr

^dInserm U869, 146 rue Léo Saignat, 33076 Bordeaux, France

†Electronic supplementary information (ESI) available. See DOI: 10.1039/c5bm00274e



Observation of the interactions between KG51 and the GTPswitch in the DNA frame

First, we examined the interaction of a pair of kissing RNA hairpins, Aptakiss and KG51, in the DNA frame (Fig. 2a). KG51 is known to be capable of binding to the Aptakiss without the need for any additional cofactors and was used as an appropriate positive control for evaluating the data using this observation system. Quantitative analysis of micrographs of this sample revealed a high percentage of X-shaped structure (84.9%), suggesting that the KG51–Aptakiss system worked well in the DNA frame.

We next substituted KG51 with the GTPswitch to examine the ligand-dependent binding property of the GTPswitch to the Aptakiss. Contrary to our expectation from the previous bulk experiment, 75.1% of DNA frame were observed to have

Table 1 Summary of the X-shape formation using Aptakiss and GTPswitch in the absence and presence of ligands

	X-shape (%)	Double-loop shape (%)	±S.D.	Counted numbers
KG51	84.9	15.1	3.3	280
GTPswitch	75.1	24.9	4.3	285

The data are represented as the mean ± S.D. of triplicate experiments ($n = 3$).

the X-shaped structure even in the absence of GTP; this percentage was only ~10% lower than that obtained for KG51 (Fig. 2c and Table 1). This unexpected high binding might have occurred because of an interaction between the bases in the kissing loop of the Aptakiss and the complementary bases of the unfolded free aptaswitch that was not previously detected in solution.¹⁶ We note that the two RNA sequences are located in relatively close positions in the DNA frame. The distance between the Aptakiss and the GTPswitch was estimated to be ~10 nm. Assuming that the motion of each aptamer is limited to a sphere with a diameter of 10 nm, the hypothetical concentration can be estimated to be ~1 mM. Although the molecular movement is constrained by fixation to the nanocavity, the molecules should behave as if they exist at a high concentration. This proximity effect may result in the ligand-independent binding.

Observation of the switching function of the GTPswitch with a cover strand

To improve the ligand dependency of the GTPswitch in the DNA frame, we used a cover strand. Such a strategy was recently demonstrated to be successful and improved the specificity of an aptamer to adenosine.¹⁶ In our case, the chosen cover strand binds to the kissing loop of the GTPswitch and extends to part of the central loop that is the GTP binding site of the aptamer (Fig. 3a). This cover strand can be displaced by the addition of GTP, which allows the GTPswitch to bind to the Aptakiss in a ligand-dependent manner. The cover strand was hybridized to the GTPswitch by annealing, and the covered GTPswitch strand was then incorporated into the DNA frame together with the strand carrying the Aptakiss. In the absence of GTP, the percentage of X-shaped structure in the AFM images was calculated as 64.4% (Fig. 3b). This value reflected a ~20% decrease in the percentage of the X-shape and is much lower than the values mentioned above for KG51 and the GTPswitch. The results indicate that the cover strand reduced the ligand-independent binding between the GTPswitch and the Aptakiss. Next, we examined the effect of introducing a cover strand-bound GTPswitch to the Aptakiss in the presence of GTP (Fig. 3c). Adenosine triphosphate (ATP) was also used to investigate the ligand specificity of the GTPswitch (Fig. 3d). From the AFM images, 82.5% of the cover strand-bound GTPswitch was in the X-shape in the presence of GTP; this value is similar to that for KG51 (84.9%). By contrast, in the presence of ATP, only 66.5% was in the X-shape; this




Fig. 2 Observations of the interactions between the Aptakiss and its counterpart either KG51 or GTPswitch in the DNA frame. (a) AFM images of the DNA frames with the Aptakiss and KG51. (b) AFM images of the DNA frames with the Aptakiss and the GTPswitch. Red and blue arrows indicate the double-loop and X-shape, respectively. Green rectangles represent an unidentified DNA frame. (c) Formation of the X-shape and double-loop in the DNA frame. Red and blue bars represent the percentages of the double-loop and X-shape formation, respectively.






Fig. 3 Observation of the interaction between the Aptakiss and the GTPswitch in the presence of a cover strand. (a) The cover strand for the GTPswitch used to prevent interaction with the counterpart Aptakiss. (b) AFM image of the DNA frames with the Aptakiss and the GTPswitch. Red and blue arrows indicate double-loop and X-shape, respectively. Green rectangles represent unidentified DNA frames. (c) AFM image of the DNA frames with the Aptakiss and the GTPswitch in the presence of GTP. (d) AFM image of the DNA frames with the Aptakiss and the GTPswitch in the presence of ATP. (e) Formation of the X-shape and double-loop in the DNA frame. Red and blue bars represent the percentages of the double-loop and X-shape, respectively. (f) Proportion of the X-shape formation in the DNA frame at various concentration of GTP and ATP.

value is close to that in the absence of GTP (64.4%). These results indicate that GTP could selectively induce binding between the GTPswitch and the Aptakiss, whereas ATP could not (Fig. 3e and Table 2).

Table 2 Summary of the X-shape formation using Aptakiss and GTPswitch with a cover strand in the absence and presence of ligands

GTPswitch/ cover strand	X-shape (%)	Double-loop shape (%)	±S.D.	Counted numbers
No ligand	64.4	35.6	4.4	381
0.1 mM GTP	74.3	25.7	1.0	238
0.5 mM GTP	79.4	20.6	0.8	242
1.0 mM GTP	82.5	17.5	2.1	324
0.1 mM ATP	67.6	32.4	0.9	216
0.5 mM ATP	67.1	32.9	1.7	229
1.0 mM ATP	66.5	33.5	2.3	281

The data are represented as the mean \pm S.D of triplicate experiments ($n = 3$).




Fig. 4 Observation of the interaction between the Aptakiss and the mutant GTPswitch in the DNA frame. (a) One G was deleted from the kissing loop of the GTPswitch to suppress the interaction with the counterpart Aptakiss. (b) AFM image of the DNA frames with the Aptakiss and the mutant GTPswitch. Red and blue arrows indicate the double-loop and X-shape, respectively. Green rectangles represent unidentified DNA frames. (c) AFM image of the DNA frames with the Aptakiss and the mutant GTPswitch in the presence of GTP. (d) AFM image of the DNA frames with the Aptakiss and the mutant GTPswitch in the presence of ATP. (e) Formation of the X-shape and double-loop in the DNA frame. Red and blue bars represent the percentages of the double-loop and X-shape formation, respectively.

Table 3 Summary of the X-shape formation using Aptakiss and GTPswitch mutant in the absence and presence of ligands

GTPswitch mutant	X-shape (%)	Double-loop shape (%)	±S.D.	Counted numbers
No ligand	44.0	56.0	2.0	202
1.0 mM GTP	65.2	34.8	0.5	230
1.0 mM ATP	46.4	53.6	2.5	214

The data are represented as the mean ± S.D of triplicate experiments ($n = 3$).

Moreover, we observed the binding in the presence of 0.1 and 0.5 mM of GTP and ATP. In the case of ATP, there is no change of the proportion. On the other hand, we found that at 0.1 mM of GTP the proportion of X-shape was decreased by ~5% from the proportion at 0.5 and 1.0 mM of GTP (Fig. 3f).

Observation of GTP switching in a mutant GTPswitch

When the Aptakiss and the GTPswitch were placed in the DNA frame, the two RNA hairpins bound easily because of the close packing in the nanospace. We tried to reduce the interaction of RNA hairpins by using a mutant GTPswitch, in which one G was deleted from the kissing loop of the GTPswitch (Fig. 4a). After assembling the Aptakiss and the mutant GTPswitch strands, we used AFM to observe the formation of the X-shape (Fig. 4b–d). In the absence of GTP, the X-shape formation between the Aptakiss and the mutant GTPswitch was 44.0%, which indicated a significant suppression of the interaction compared with the X-shape formed with the usual GTPswitch (75.1%). To examine the switching ability, formation of the X-shape was observed in the presence of GTP. The percentage in the X-shape was observed as 65.2%, which indicated a 21% increase in the binding of the GTPswitch and the Aptakiss when GTP was added. To confirm the ligand selectivity, we added ATP instead of GTP. In the presence of ATP, the percentage of the X-shape decreased by 19% to be 46.4% compared with that observed in the presence of GTP (Fig. 4e and Table 3). These results indicate that the mutant GTPswitch preserved the switching ability and ligand selectivity. These findings indicate that adjusting the association and dissociation of the kissing interaction of the RNA hairpins by deleting the nucleotide in the kissing domain was successful without losing the switching ability.

Conclusions

We performed single-molecule observations of kissing complexes in the nanocavity of the DNA origami frame. Intriguingly, in the closely spaced condition, the GTPswitch could bind to the Aptakiss even in the absence of GTP in contrast to previous work.^{13,16} This GTP-independent binding could be suppressed by the addition of a cover strand against the kissing loop of the GTPswitch, and the GTP-dependent binding of the GTPswitch and Aptakiss was observed. The

mutant GTPswitch also worked to control the kissing interaction and exhibited preserved switching ability and ligand selectivity. Although further optimization in the switching response is required, these findings support the potential applications of ligand-responsive kissing aptamers for dynamic systems that can be organized on DNA origami nanostructures. We believe that ligand-responsive kissing aptamers will enable us to regulate more global changes in nucleic acid nanostructures, such as programmed oligomerization into prescribed patterns.

Acknowledgements

This work was supported by JSPS KAKENHI (grant numbers 15H03837, 24104002, 24225005, 26620133) to ME and HS. Financial supports from the Sekisui Chemical Research Grant and the Kurata Memorial Hitachi Science and Technology Foundation to ME are also acknowledged. We acknowledge the financial support of the Aquitaine Regional Government and ANR program VIBBnano (ANR-10-NANO-04) to JJT (Bordeaux).

Notes and references

- 1 F. Beaurain, C. Di Primo, J. J. Toulmé and M. Laguerre, *Nucleic Acids Res.*, 2003, **31**, 4275–4284.
- 2 I. Lebars, P. Legrand, A. Aime, N. Pinaud, S. Fribourg and C. Di Primo, *Nucleic Acids Res.*, 2008, **36**, 7146–7156.
- 3 J. C. Paillart, E. Skripkin, B. Ehresmann, C. Ehresmann and R. Marquet, *Proc. Natl. Acad. Sci. U. S. A.*, 1996, **93**, 5572–5577.
- 4 J. I. Tomizawa, *Cell*, 1986, **47**, 89–97.
- 5 C. Persson, E. G. H. Wagner and K. Nordström, *EMBO J.*, 1990, **9**, 3761–3775.
- 6 L. Argaman and S. Altuvia, *J. Mol. Biol.*, 2000, **300**, 1101–1112.
- 7 C. Brunel, R. Marquet, P. Romby and C. Ehresmann, *Biochimie*, 2002, **84**, 925–944.
- 8 E. Goux, S. Lisi, C. Ravelet, G. Durand, E. Fiore, E. Dausse, J. J. Toulmé and E. Peyrin, *Anal. Bioanal. Chem.*, 2015, **407**, 6515–6524.
- 9 H. Van Melckebeke, M. Devany, C. Di Primo, F. Beaurain, J. J. Toulmé, D. L. Bryce and J. Boissbouvier, *Proc. Natl. Acad. Sci. U. S. A.*, 2008, **105**, 9210–9215.
- 10 W. W. Grabow and L. Jaeger, *Acc. Chem. Res.*, 2014, **47**, 1871–1880.
- 11 A. Chworos, I. Severcan, A. Y. Koyfman, P. Weinkam, E. Oroudjev, H. G. Hansma and L. Jaeger, *Science*, 2004, **306**, 2068–2072.
- 12 P. Guo, *Nat. Nanotechnol.*, 2010, **5**, 833–842.
- 13 F. Ducongé and J. J. Toulmé, *RNA*, 1999, **5**, 1605–1614.
- 14 K. Kikuchi, T. Umehara, K. Fukuda, J. Hwang, A. Kuno, T. Hasegawa and S. Nishikawa, *J. Biochem.*, 2003, **133**, 263–270.



- 15 S. Da Rocha Gomes, E. Dausse and J. J. Toulmé, *Biochem. Biophys. Res. Commun.*, 2004, **322**, 820–826.
- 16 G. Durand, S. Lisi, C. Ravelet, E. Dausse, E. Peyrin and J. J. Toulmé, *Angew. Chem., Int. Ed.*, 2014, **53**, 6942–6945.
- 17 M. Endo and H. Sugiyama, *Acc. Chem. Res.*, 2014, **47**, 1645–1653.
- 18 A. Rajendran, M. Endo and H. Sugiyama, *Chem. Rev.*, 2014, **114**, 1493–1520.
- 19 (a) Y. Sannohe, M. Endo, Y. Katsuda, K. Hidaka and H. Sugiyama, *J. Am. Chem. Soc.*, 2010, **132**, 16311–16313; (b) A. Rajendran, M. Endo, K. Hidaka and H. Sugiyama, *Angew. Chem., Int. Ed.*, 2014, **53**, 4107–4112.
- 20 (a) A. Rajendran, M. Endo, K. Hidaka and H. Sugiyama, *J. Am. Chem. Soc.*, 2013, **135**, 1117–1123; (b) M. Endo, M. Inoue, Y. Suzuki, C. Masui, H. Morinaga, K. Hidaka and H. Sugiyama, *Chem. – Eur. J.*, 2013, **19**, 16887–16890.
- 21 (a) M. Endo, Y. Yang, Y. Suzuki, K. Hidaka and H. Sugiyama, *Angew. Chem., Int. Ed.*, 2012, **51**, 10518–10522; (b) Y. Yang, M. Endo, Y. Suzuki, K. Hidaka and H. Sugiyama, *Chem. Commun.*, 2014, **50**, 4211–4213.
- 22 (a) M. Endo, Y. Katsuda, K. Hidaka and H. Sugiyama, *J. Am. Chem. Soc.*, 2010, **132**, 1592–1597; (b) M. Endo, Y. Katsuda, K. Hidaka and H. Sugiyama, *Angew. Chem., Int. Ed.*, 2010, **49**, 9412–9416; (c) Y. Suzuki, M. Endo, Y. Katsuda, K. Ou, K. Hidaka and H. Sugiyama, *J. Am. Chem. Soc.*, 2014, **136**, 211–218; (d) Y. Suzuki, M. Endo, C. Cañas, S. Ayora, J. C. Alonso, H. Sugiyama and K. Takeyasu, *Nucleic Acids Res.*, 2014, **42**, 7421–7428.

